Astronomy & Astrophysics manuscript no. Faifer˙et˙al
February 22, 2017
© ESO 2017

arXiv:1702.06472v1 [astro-ph.GA] 21 Feb 2017

Letter to the Editor

First confirmed ultra-compact dwarf galaxy in the NGC 5044 group

Favio R. Faifer[1,2], Carlos G. Escudero[1,2], María C. Scalia[1,2], Analía V. Smith Castelli[2,3], Mark Norris[4], María E. de Rossi[5,6], Juan C. Forte[3,7], and Sergio A. Cellone[1,2]

[1] Facultad de Ciencias Astronómicas y Geofísicas, Universidad Nacional de La Plata, Paseo del Bosque s/n, B1900FWA, La Plata, Argentina
e-mail: favio@fcaglp.unlp.edu.ar
[2] Instituto de Astrofísica de La Plata (CCT La Plata - CONICET - UNLP), Paseo del Bosque s/n, B1900FWA, La Plata, Argentina
[3] Consejo Nacional de Investigaciones Científicas y Técnicas, Godoy Cruz 2290, C1425FQB, CABA, Argentina
[4] Jeremiah Horrocks Institute, University of Central Lancashire, Preston, PR1 2HE, United Kingdom
[5] Universidad de Buenos Aires, Facultad de Ciencias Exactas y Naturales y Ciclo Bsico Comn. Buenos Aires, Argentina
[6] CONICET-Universidad de Buenos Aires, Instituto de Astronoma y Fsica del Espacio (IAFE).
[7] Planetario 'Galileo Galilei', Secretaría de Cultura, Av. Sarmiento S/N, 1425 Ciudad Autnoma de Buenos Aires, Argentina

Received March 1, 2017; accepted March 16, 2017

ABSTRACT

Context. Ultra-compact dwarfs (UCDs) are stellar systems displaying colours and metallicities between those of globular clusters (GCs) and early-type dwarf galaxies, as well as sizes of $R_{\rm eff} \lesssim 100$ pc and luminosities in the range $-13.5 < M_{\rm V} < -11$ mag. Although their origin is still subject of debate, the most popular scenarios suggest that they are massive star clusters or the nuclei of tidally stripped dwarf galaxies.
Aims. NGC 5044 is the central massive elliptical galaxy of the NGC 5044 group. Its GC/UCD system is completely unexplored.
Methods. In Gemini+GMOS deep images of several fields around NGC 5044 and in spectroscopic multi-object data of one of these fields, we detected an unresolved source with $g' \sim 20.6$ mag, compatible with being an UCD. Its radial velocity was obtained with FXCOR and the penalized pixel-fitting (pPXF) code. To study its stellar population content, we measured the Lick/IDS indices and compared them with predictions of single stellar population models, and we used the full spectral fitting technique.
Results. The spectroscopic analysis of the UCD revealed a radial velocity that agrees with the velocity of the elliptical galaxy NGC 5044. From the Lick/IDS indices, we have obtained a luminosity-weighted age and metallicity of $11.7^{+1.4}_{-1.2}$ Gyr and $[Z/H] = -0.79 \pm 0.04$ dex, respectively, as well as $[\alpha/Fe] = 0.30 \pm 0.06$. From the full spectral fitting technique, we measured a lower age (8.52 Gyr) and a similar total metallicity ($[Z/H] = -0.86$ dex).
Conclusions. Our results indicate that NGC 5044-UCD1 is most likely an extreme GC ($M_V \sim -12.5$ mag) belonging to the GC system of the elliptical galaxy NGC 5044.

Key words. galaxies: elliptical and lenticular – galaxies: photometry – galaxies: star clusters: general

1. Introduction

Ultra-compact dwarfs (UCDs) are stellar systems displaying masses, colours, and metallicities within the range covered by those of globular clusters (GCs) and dwarf galaxies. They show typical effective radii ($R_{\rm eff}$) $\lesssim$100 pc and luminosities in the range $-13.5 < M_{\rm V} < -11$ mag (Mieske 2006). It now seems likely that they are a mixed population of massive star clusters and the nuclei of tidally stripped dwarf galaxies (i.e. Hilker 2009; Brodie et al. 2011; Norris & Kannappan 2011; Pfeffer et al. 2014; Longmore et al. 2014; Zhang et al. 2015). However, it has so far proved difficult to conclusively demonstrate that any individual UCD is of either type (and in particular, that it is of massive star cluster type).

Initially, UCDs were identified in the environments of rich galaxy clusters like Virgo and Fornax. However, more recently, several authors have reported UCDs in groups of galaxies and associated with galaxies in the field (Evstigneeva et al. 2007; Hau et al. 2009; Norris & Kannappan 2011; Norris et al. 2014). The existence of such objects in low-density environments provides an important insight into the relative frequency of the different formation pathways. However, deep spectroscopy, which gives information about the stellar population content among other properties, is still scarce (Francis et al. 2012; Janz et al. 2016).

We present the spectroscopic confirmation of the first UCD in the vicinity of the massive elliptical galaxy NGC 5044 ($M_{\rm B} = -21.2$ mag, $D = 35.7 \pm 5$ Mpc, $V_{\rm r} = 2782$ km s^{-1}), the central object of the NGC 5044 group. This group harbours $\sim$ 150 members, most of which are dwarf galaxies (Ferguson & Sandage 1990). This population represents half the members of the Fornax Cluster. However, its central density and velocity dispersion are higher, while its kinematics and early-type galaxy fraction reveal a mature group (Cellone & Buzzoni 2005). Although the elliptical galaxy NGC 5044 has been the subject of several studies, its rich GC system ($\sim 5500 \pm 500$ members, Faifer et al. in preparation) remains completely unexplored.

Through the analysis of Gemini+GMOS images and spectra, we have been able to obtain the photometric parameters and the radial velocity of the UCD, as well as information about its stellar population content. The letter is organized as follows. In

Table 1. Log of the GEMINI-GMOS data.

Date	Filter/Grism	Exposures (sec)	FWHM/Resolution (arcsec/Å)
27 Jan. 2009	g'	11×450	0.92
	r'	2×350	0.76
	i'	7×350	0.80
26 Feb. 2009	g'	4×450	
	r'	2×350	
22 Mar. 2010	B600_G5323	3×2330	4.61
19 Apr. 2010	B600_G5323	3×2330	4.61
20 Apr. 2010	B600_G5323	2×2330	4.61

Section 2 we present our data, in Section 3 our analysis and results, and in Section 4 we discuss the results and we present the conclusions.

2. Observational data

With the Gemini Multi-Object Spectrograph (GMOS) (Hook et al. 2004) of GEMINI-South, we obtained deep g', r', and i' images of several fields around NGC 5044 (Program: GS-2009A-Q-46). In this letter we focus on the field that contains the elliptical galaxy. The analysis of the complete photometric dataset will be presented in a forthcoming paper (Faifer et al. in preparation). GMOS consists of three CCDs of 2048 × 4096 pixels, separated by gaps of ∼ 2.8 arcsec, with an unbinned pixel scale of 0.0727 arcsec pixel^{-1}. The field of view (FOV) is 5.5' × 5.5', and the scale for binning 2×2 is ∼ 0.146 arcsec pixel^{-1}. The images were reduced in the usual way (Escudero et al. 2015).

We have also obtained a GMOS spectroscopic mask of the field containing NGC 5044 with the aim at obtaining radial velocities of a sample of GC candidates (program: GS-2010A-Q-56). The instrumental setup included 2×2 binning and slits of 1″ width, with dithering in the spectral direction in order to fill the CCD gaps. The data were reduced using the IRAF tasks gbias, gsflat, gsreduce, and mgswavelength. The individual spectra were then extracted using apall and were combinded through the task scombine. Finally, an approximate flux calibration was applied based on the flux of the standard star LTT 4816. The final spectrum covers a spectral range of 3750-6780 Å and displays a resolution of ∼ 4.61 Å. The signal-to-noise ratio (S/N) per Å of the final spectrum increases from 20 at 4100 Å to 70 at 6000 Å. In Table 1 we show the basic information related with all our data.

3. Analysis and results

3.1. Photometry

We have subtracted the galaxy light from our images using an iterative combination of the background-modelling tool of SExtractor (Bertin & Arnouts 2010) and a median filtering, following Puzia et al. (2004). SExtractor was also used to perform the detection of GC candidates in the field. At the distance adopted here for NGC 5044, 1 arcsec is equivalent to 170 pc. This means that any globular cluster or UCD present in the field will appear as an unresolved source in our images. We have used Daophot to obtain the photometry of the sources detected in the field, and we modelled the psf using ∼ 25-30 well-exposed unresolved objects.

Figure 1 shows the colour-magnitude diagram for the unresolved sources detected in the field. The magnitudes and colours were calibrated to the Sloan Digital Sky Survey (SDSS) photometric system and corrected for interstellar extinction (Schlafly & Finkbeiner 2011). Objects with integrated colours in the ranges 0.5< $(g'-i')_0$ <1.4, 0.35< $(g'-r')_0$ <0.85 and 0.0< $(r'-i')_0$ <0.55 and magnitudes $g'_0 > 22$ were considered as GC candidates (Faifer et al. 2011). Three UCD candidates were selected using the same colour ranges, but with $g'_0 < 22$, which translates into $M_V < -11$ mag at the NGC 5044 distance (Mieske et al. 2006).

Table 2. Basic information of NGC 5044-UCD1. Apparent magnitudes and colours are not extinction or reddening corrected.

R.A. (J2000)	13:15:35.75	
DEC (J2000)	-16:23:25.33	
A_g (mag)	0.231	
A_r (mag)	0.160	
A_i (mag)	0.119	
$m_{g'}$ (mag)	20.62	± 0.01
$(g'-r')$ (mag)	0.67	± 0.01
$(g'-i')$ (mag)	0.98	± 0.01
$(r'-i')$ (mag)	0.31	± 0.01
Vr (km s^{-1})	2945 ± 5.00	
Age_{Lick} (Gyr)	11.7	± 1.4
$[Z/H]_{Lick}$	-0.79	± 0.04
$[\alpha/Fe]$	0.30	± 0.06
M_{ssp} ($M_\odot$)	2.8×10^7	$\pm 3 \times 10^6$
Age_{pPXF} (Gyr)	8.52	± —
$[Z/H]_{pPXF}$	-0.86	± —
H_{δ_A} (Å)	1.46	± 0.36
H_{δ_F} (Å)	1.53	± 0.24
G_{4300} (Å)	3.59	± 0.28
Fe_{4383} (Å)	2.47	± 0.36
H_β (Å)	2.18	± 0.12
Fe_{5015} (Å)	3.01	± 0.25
Mgb (Å)	2.35	± 0.11
Fe_{5270} (Å)	1.68	± 0.13
Fe_{5335} (Å)	1.69	± 0.14
Fe_{5406} (Å)	0.74	± 0.11
$[MgFe]'$ (Å)	1.99	± 0.11
$\langle Fe \rangle$ (Å)	1.69	± 0.10

3.2. Radial velocity

A preliminary estimation of the radial velocity (V_r) of the three UCD candidates was obtained from the spectroscopic data using the task FXCOR from IRAF. As spectral templates, we used the stellar population synthesis models MILES (Vazdekis et al. 2010) considering two total metallicities ($[Z/H] = -1.31$ and $[Z/H] = 0.00$ dex), a bimodal IMF with a slope of 1.3, and ages of 12 Gyr. For two of these objects we have obtained $V_r < 400$ km s^{-1}, meaning that they are Milky Way (MW) stars. For the third object (hereafter, NGC 5044-UCD1) we have obtained $V_r \sim 2900$ km s^{-1}, similar to the reported systemic velocity of the galaxy NGC 5044 (2782 km s^{-1}), thereby confirming that NGC 5044-UCD1 is a member of this system.

Subsequently, we used the penalized pixel-fitting code pPXF (Cappellari & Emsellem 2004) in order to mitigate the mismatch template effect and, therefore, to better estimate the V_r of the UCD. We used the SSP MILES library models from Vazdekis et al. (2010), which span a range in age of 0.03-14 Gyr and $-2.27 < [Z/H] < 0.4$ dex. We obtained a radial velocity of $V_r \sim 2945$ km s^{-1}. This value and the error quoted in Table 2 were obtained as the median from 500 Monte Carlo real-



Fig. 1. Colour-magnitude diagram of the unresolved sources detected in the field of NGC 5044 (dots). Small circles depict the objects that match our colour selection for globular clusters. The small circles with a cross inside show two candidates whose radial velocities proved that they are MW stars. The circle surrounded by the ring shows the location of NGC 5044-UCD1 in the diagram.



Fig. 2. Integrated spectrum of NGC 5044-UCD1 (black) and one of the kinematics fits obtained by pPXF (grey). We also show the residuals of the fit offset by 0.5, for clarity.

izations and the 1σ estimation, respectively. Figure 2 shows the integrated spectrum of the UCD together with an example of the fits obtained by pPXF.

3.3. Age and metalicity from Lick indices and SSP models

We have obtained the Lick indices from the GMOS spectra following Norris et al. (2008, 2015) (Table 2). Uncertainties for the indices were obtained from 500 Monte Carlo simulations based on a combination of errors introduced by photon noise (both target and sky) and errors in the measured redshift of the spectra (essentially negligible here). The age, metallicity, and α-elements ratio of NGC 5044-UCD1 were obtained through the χ^2 minimization method of Proctor & Sansom (2002) and Proctor et al. (2004). We have used a smooth grid in age, [α/Fe], and [Z/H] by interpolating the single stellar population (SSP) models of Thomas et al. (2003, 2004). As in similar works (e.g. Norris et al. 2006), the indices $H_{\delta A}$, $H_{\delta F}$, $H\gamma A$ G4300, Fe4383,



Fig. 3. Diagnostic diagram H_β vs $[MgFe]$. We also overplot the models of Thomas et al. (2003, 2004) with $[\alpha/Fe] = 0.3$. NGC 5044-UCD1 (filled circle with error bars) is an ancient object with subsolar metallicity.

$H\beta$, Fe5015, Mgb, Fe5270, Fe5335, and Fe5406 gave the most reliable results. The analysis provides an age of $11.7^{+1.4}_{-1.2}$ Gyr, a total metallicity of $[Z/H] = -0.79 \pm 0.04$ dex, and an α-elements ratio of $[\alpha/Fe] = 0.30 \pm 0.06$. From these values and using the M/L ratios given by Maraston (2005) for a Kroupa IMF (M/L = 2.9 ± 0.3), we infer a total stellar mass of $2.8(\pm 0.3) \times 10^7\ M_\odot$. In Figure 3 we show the diagnostic diagram H_β vs $[MgFe]$, where $[MgFe]' = [Mgb \times (0.72 \times Fe5270 + 0.28 \times Fe5335)]^{0.5}$, overplotting the models of Thomas et al. (2003, 2004) with $[\alpha/Fe] = 0.3$, according to the value previously estimated.

3.4. Star formation history

Through the full spectral fitting technique implemented within the pPXF code, we studied the star formation history (SFH) of NGC 5044-UCD1 and obtained new estimates of the age and metallicity of this object in a way that is independent of that presented in Section 3.3. We applied the regularization method to obtain a smooth solution of the best fit linear combination of SSP models to the UCD spectrum (McDermid et al. 2015). We used the same library of models presented in Section 3.2, adopting an α-element ratio of $[\alpha/Fe] = 0.4$, which is close to the value obtained in the Lick index analysis.

We performed several pPXF runs considering different spectral ranges of the UCD spectrum. In all the cases, we achieved a smooth solution that included a main contribution of models around a mean age of 8.52 Gyr and a mean metalicity of $[Z/H] = -0.86$ dex (Figure 4). In some of the tests, which include the regions of the highest S/N ($\lambda > 4500$Å), we also obtained a secondary contribution of models with higher metallicities ($[Z/H] > -0.25$ dex). In Figure 4 we show one of the tests with both contributions. To check this result, we built a synthetic spectrum combining two SSP models with the ages and metallicities obtained from our analysis for the two components. We added to this spectrum a noise level similar to that found in the observed spectrum, and we ran pPFX. We recovered the original two components, but a third spurious subpopulation was also



Fig. 4. Star formation history of NGC 5044-UCD1 obtained using pPXF. The different grey-scale regions indicate the weight of each SSP model, which is equivalent to the zero-age mass distribution of the SFH. The dashed lines indicate the $[Z/H]$ of each available model. The black filled circle depicts the luminosity-weighted SSP age and metallicity of this object as measured using absorption line strength indices (Section 3.3). The white pentagon shows the location of the mass-weighted age and metallicity derived from the full spectral fitting technique, and considering only models with $[Z/H] < -0.25$ dex.

obtained. To confirm this result, we would need a spectrum displaying a higher S/N in the whole spectral range.

4. Discussion and conclusions

From the photometric and spectroscopic analysis of a deep field taken with Gemini+GMOS, we have been able to detect and confirm the first UCD in the NGC 5044 group. Its radial velocity and angular proximity (2.83 arcmin) indicate that this object is associated with the massive early-type galaxy NGC 5044. At the distance adopted in this work for this galaxy, this UCD appears to be an unresolved object, and it has an absolute magnitude of $M_V = -12.5$ mag. This luminosity is well above the usual upper cut for "classical" globular clusters (Faifer et al 2011).

We studied its stellar population properties using two methods: the analysis of Lick/IDS indices, and the application of the full spectral fitting technique. We find a discrepancy in the age determined by the two methods. From the comparison of the Lick/IDS values with SSP models, we have obtained an age of $11.7^{+1.4}_{-1.2}$ Gyr, which is significantly older than the value of 8.5 Gyr obtained with the full spectral fitting technique, although the uncertainties on the full spectral fitting age are difficult to quantify. Similar discrepancies have been reported for UCDs before (e.g. Francis et al. 2012). However, in contrast to most of the objects presented by Francis and collaborators, our metallicity values are consistent regardless of the method used ($[Z/H] = -0.79 \pm 0.04$ dex and $[Z/H] = -0.86$ dex). It is notable that the metallicity of NGC 5044-UCD1 is within the range displayed by the UCDs detected in Virgo and Fornax, but considerably lower than that of the confirmed stripped nuclei presented by Janz et al. (2016).

In common with most globular clusters (Puzia et al. 2005; Norris et al. 2008), NGC 5044-UCD1 presents a supersolar α-element abundance of $[\alpha/Fe] = 0.30$, suggesting rapid star formation. The SFH analysis suggests the presence of a secondary metal-rich stellar population, of an age indistinguishable from the main population. If confirmed by higher S/N data, this result could be associated with the "blue tilt" phenomena of massive GCs, whereby the most massive low-metallicity GCs are found to be progressively redder. This has been interpreted as being due to self-enrichment, where the most massive GCs are able to form additional generations of stars enriched in metals from the ejecta of the initial population (Strader & Smith 2008).

According to Francis et al. (2012), the UCDs of Virgo and Fornax do not display properties similar to those of the nuclei of present-day nucleated dwarf galaxies. This picture seems to be in agreement with the location that our UCD would have in Figure 13 of Norris & Kannappan (2011). In that figure, the absolute visual magnitude of our object ($M_V = -12.5$ mag) and the total number of GCs of NGC 5044 ($\sim 5500 \pm 500$, Faifer et al. in preparation) would agree with the general trend defined by the UCDs that are more likely the brightest tail of the GC population. In this sense, from a Monte Carlo experiment using the luminosity function of the NGC 5044 GC system, we have obtained that in 30% of the realizations we have $1-3$ GCs brighter than $M_V = -12.5$ mag. Therefore, all the lines of evidence would point toward NGC 5044-UCD1 being an unusually massive GC of the NGC 5044 system.

Acknowledgements. Based on observations obtained at the Gemini Observatory (Programs GS-2009A-Q-46 and GS-2010A-Q-56), which is operated by the Association of Universities for Research in Astronomy, Inc., under a cooperative agreement with the NSF on behalf of the Gemini partnership: the National Science Foundation (United States), the National Research Council (Canada), CONICYT (Chile), the Australian Research Council (Australia), Ministério da Ciência, Tecnologia e Inovação (Brazil) and Ministerio de Ciencia, Tecnología e Innovación Productiva (Argentina).

This research has made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

This work was funded with grants from Consejo Nacional de Investigaciones Científicas y Técnicas and Universidad Nacional de La Plata (Argentina).

References

Bertin, E. & Arnouts, S. 2010, SExtractor: Source Extractor, Astrophysics Source Code Library
Brodie, J. P., Romanowsky, A. J., Strader, J., & Forbes, D. A. 2011, AJ, 142, 199
Cappellari, M. & Emsellem, E. 2004, PASP, 116, 138
Cellone, S. A. & Buzzoni, A. 2005, MNRAS, 356, 41
Escudero, C. G., Faifer, F. R., Bassino, L. P., Calderón, J. P., & Caso, J. P. 2015, MNRAS, 449, 612
Evstigneeva, E. A., Drinkwater, M. J., Jurek, R., et al. 2007, MNRAS, 378, 1036
Faifer, F. R., Forte, J. C., Norris, M. A., et al. 2011, MNRAS, 416, 155
Ferguson, H. C. & Sandage, A. 1990, AJ, 100, 1
Francis, K. J., Drinkwater, M. J., Chilingarian, I. V., Bolt, A. M., & Firth, P. 2012, MNRAS, 425, 325
Hau, G. K. T., Spitler, L. R., Forbes, D. A., et al. 2009, MNRAS, 394, L97
Hilker, M. 2009, in Reviews in Modern Astronomy, Vol. 21, Reviews in Modern Astronomy, ed. S. Röser, 199–213
Hook, I. M., Jørgensen, I., Allington-Smith, J. R., et al. 2004, PASP, 116, 425
Janz, J., Norris, M. A., Forbes, D. A., et al. 2016, MNRAS, 456, 617
Longmore, S. N., Kruijssen, J. M. D., Bastian, N., et al. 2014, Protostars and Planets VI, 291
Maraston, C. 2005, MNRAS, 362, 799
McDermid, R. M., Alatalo, K., Blitz, L., et al. 2015, MNRAS, 448, 3484
Mieske, S. 2006, in Revista Mexicana de Astronomia y Astrofisica Conference Series, Vol. 26, Revista Mexicana de Astronomia y Astrofisica Conference Series, 194
Mieske, S., Hilker, M., Infante, L., & Jordán, A. 2006, AJ, 131, 2442
Norris, M. A., Escudero, C. G., Faifer, F. R., et al. 2015, MNRAS, 451, 3615

Norris, M. A. & Kannappan, S. J. 2011, MNRAS, 414, 739
Norris, M. A., Kannappan, S. J., Forbes, D. A., et al. 2014, MNRAS, 443, 1151
Norris, M. A., Sharples, R. M., Bridges, T., et al. 2008, MNRAS, 385, 40
Norris, M. A., Sharples, R. M., & Kuntschner, H. 2006, MNRAS, 367, 815
Pfeffer, J., Griffen, B. F., Baumgardt, H., & Hilker, M. 2014, MNRAS, 444, 3670
Proctor, R. N., Forbes, D. A., & Beasley, M. A. 2004, MNRAS, 355, 1327
Proctor, R. N. & Sansom, A. E. 2002, MNRAS, 333, 517
Puzia, T. H., Kissler-Patig, M., Thomas, D., et al. 2005, A&A, 439, 997
Puzia, T. H., Kissler-Patig, M., Thomas, D., et al. 2004, A&A, 415, 123
Schlafly, E. F. & Finkbeiner, D. P. 2011, ApJ, 737, 103
Strader, J. & Smith, G. H. 2008, AJ, 136, 1828
Thomas, D., Maraston, C., & Bender, R. 2003, MNRAS, 339, 897
Thomas, D., Maraston, C., & Korn, A. 2004, MNRAS, 351, L19
Vazdekis, A., Sánchez-Blázquez, P., Falcón-Barroso, J., et al. 2010, MNRAS, 404, 1639
Zhang, H.-X., Peng, E. W., Côté, P., et al. 2015, ApJ, 802, 30